



### Synonym

Integrin alpha 8 beta 1, ITGA8&ITGB1

### Source

Rat ITGA8 & ITGB1 Heterodimer Protein, His Tag&Tag Free (IT1-R53W6) is expressed from CHO cells. It contains AA Phe 38 - Pro 1034 (ITGA8) & Gln 21 - Asp 729 (ITGB1) (Accession # [F7F1E3](#) (ITGA8) & [P49134](#) (ITGB1)).

Predicted N-terminus: Phe 38 (ITGA8) & Gln 21 (ITGB1)

### Molecular Characterization

ITGA8 (Phe 38 - Pro 1034) F7F1E3	Acidic Tail	His
ITGB1 (Gln 21 - Asp 729) P49134	Basic Tail	

Rat ITGA8 & ITGB1 Heterodimer Protein, His Tag&Tag Free, produced by co-expression of ITGA8 and ITGB1, has a calculated MW of 117.2 kDa (ITGA8) and 83.7 kDa (ITGB1). The protein migrates as 50-55 kDa and 65-150 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under non-reducing (NR) condition (SDS-PAGE) due to glycosylation.

### Endotoxin

Less than 1.0 EU per µg by the LAL method.

### Purity

>80% as determined by SDS-PAGE.

### Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 150 mM NaCl, pH 7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

### Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

### Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

### SDS-PAGE



Rat ITGA8 & ITGB1 Heterodimer Protein, His Tag&Tag Free on SDS-PAGE under non-reducing (NR) condition. The gel was stained with Coomassie Blue.

The purity of the protein is greater than 80% (With [Star Ribbon Pre-stained Protein Marker](#)).

### Background

Human Integrin alpha 8 beta 1 Heterodimer Protein consists of ITGA8 and ITGB1. The integrin alpha 8 subunit, isolated by low stringency hybridization, is a novel integrin subunit that associates with beta 1. The recently identified alpha 8 integrin subunit associates with beta 1 and is predominantly expressed in smooth muscle

Discounts, Gifts,  
and more!



# Rat ITGA8 & ITGB1 Heterodimer Protein, His Tag&Tag Free

Catalog # IT1-R53W6



and other contractile cells in adult tissues, and in mesenchymal and neural cells during development. In addition, Integrin alpha 8 beta 1 is a receptor for fibronectin and can promote attachment, cell spreading, and neurite outgrowth on fibronectin.

## Clinical and Translational Updates

Please contact us via [TechSupport@acrobiosystems.com](mailto:TechSupport@acrobiosystems.com) if you have any question on this product.

Discounts, Gifts,  
and more!



➤ [www.acrobiosystems.com](http://www.acrobiosystems.com)

12/26/2023