


Synonym

CO-029,tetraspanin 8, TM4SF3, TM4SF3tspan-8, Transmembrane 4 superfamily member 3tetraspanin-8, TSPAN8, Tspan-8, Tumor-associated antigen CO-029

Source

Mouse TSPAN8 Protein, His Tag(TS8-M5249) is expressed from human 293 cells (HEK293). It contains AA Gly 106 - Asn 203 (Accession # [Q8R3G9-1](#)). Predicted N-terminus: HiS

Molecular Characterization



This protein carries a polyhistidine tag at the N-terminus

The protein has a calculated MW of 12.8 kDa. The protein migrates as 15-19 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Mouse TSPAN8 Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.

SEC-MALS



The purity of Mouse TSPAN8 Protein, His Tag (Cat. No. TS8-M5249) is more than 90% and the molecular weight of this protein is around 13-18 kDa verified by SEC-MALS.

[Report](#)

Background

Tspan8 is 1 of the 33 mammalian members of the tetraspanin family, composed of transmembrane proteins that organize laterally, together or with other membrane partners such as integrins, to form ‘tetraspanin webs’. These platforms signal within cells to regulate many cellular processes: adhesion, migration, invasion or survival Tspan8 has been implicated in many types of cancer. Overexpression was reported in glioma and colorectal, esophageal, hepatic, gastric and pancreatic

carcinoma. Tspan8 exerts a pro-invasive function by controlling cell–cell and cell–matrix interactions through its association with membrane partners such as $\alpha\beta4$ integrin-protein kinase C (PKC)-activated, E-cadherin, EpCAM, claudin-7 and CD44. Moreover, Tspan8 may be a promising new therapeutic target, as Tspan8-specific antibodies were shown to reduce cell motility, block tumor angiogenesis in vivo.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.